

**Ecologically Significant Wetlands
in the
Upper Yellowstone River Watershed**

**including the
Boulder, Clarks Fork Yellowstone,
Shields, and Stillwater River Drainages**

Prepared for the
Montana Department of Environmental Quality

By
W. Marc Jones

August 2001



**Ecologically Significant Wetlands
in the
Upper Yellowstone River Watershed

including the
Boulder, Clarks Fork Yellowstone,
Shields, and Stillwater River Drainages**

Prepared for the
Montana Department of Environmental Quality
Agreement # 200102

By
W. Marc Jones

Montana Natural Heritage Program
Montana State Library
P.O. Box 201800
Helena, Montana 59620-1800

© 2001 Montana Natural Heritage Program

This document should be cited as:

Jones, W. M. 2001. Ecologically significant wetlands in the upper Yellowstone River watershed, including the Boulder, Clarks Fork Yellowstone, Shields, and Stillwater River drainages. Report to the Montana Department of Environmental Quality. Montana Natural Heritage Program, Helena, MT. 37pp. plus appendices.

Summary

The Montana Natural Heritage Program, in partnership with the Montana Department of Environmental Quality, has completed an inventory of ecologically significant wetlands in the watersheds of the upper Yellowstone River in south central Montana. This project identified high quality wetlands in the study area and evaluated their diversity and integrity. Building on previous watershed inventories, this work creates a consistent and comprehensive source of wetland information that can form the basis for effective prioritization of wetland protection and restoration efforts.

This inventory targeted wetlands with intact hydrological functions, representative native plant communities, outstanding wildlife values, and/or rare plant and animal species. Inventory priorities were also influenced by degree of threat. Therefore highly protected alpine wetlands in the Absaroka-Beartooth Wilderness Area and the proposed Line Creek Plateau Research Natural Area were not inventoried, despite the ecological importance of these wetlands. Instead, greater priority was placed on inventorying wetlands on private land because of the greater development potential at these sites. Important sources for locating significant wetlands were local expert opinion and aerial photographs.

We used standard Heritage Program methodologies to inventory wetlands and to assess site condition, catalog community types, and document rare plant and animal occurrences. Five criteria were used to evaluate each site's ecological significance: (1) condition, which includes degree of hydrologic or geomorphic alteration, quality of native plant communities, and presence of exotic species, (2) landscape context, which includes condition of uplands and hydrologic connectivity between wetland and uplands, (3) diversity, which includes the number of plant communities, structural vegetation types,

and hydrologic classes, (4) rarity, which includes the number and condition of rare plants, animals, or communities, and (5) size of wetland. We then placed sites into one of four categories, ranging from highest quality (A-ranked) to poorest quality (D-ranked).

Forty-six ecologically significant wetlands were inventoried for this study. Of these sites, eight rated as A-ranked wetlands, 16 as B-ranked wetlands, 20 as C-ranked wetlands, and two sites were not ranked. A-ranked sites were relatively undisturbed to pristine. In general, their natural hydrologic regimes were intact, they supported high quality examples of native plant communities, and they had no or only minor weed populations. The uplands surrounding these sites were largely undisturbed, with minimal human alterations. These wetlands included diverse beaver-influenced wetlands and several poor fens, which are a regionally rare wetland type. In contrast, B-ranked sites had been impacted by both on- and off-site human disturbances, although many sites still maintained high functional capacity and supported high quality plant communities. This category included riverine and depressional montane wetlands along the Beartooth Front and low-elevation riverine and slope wetlands. Grove Creek Aspens, a unique spring-fed aspen stand in the arid Bighorn Basin, was included in this category. The remaining sites were rated as C-ranked wetlands. These wetlands have been functionally impaired through hydrologic or geomorphic alterations or through land use disturbances in the wetlands or adjacent uplands. Exotic species were widespread and abundant at many of these sites. In contrast, some of these wetlands were in good condition, but were comprised of a few common, structurally simple communities, and therefore had low diversity and rarity scores. C-ranked sites included low-elevation riverine wetlands as well as three large alkaline lake systems.

Acknowledgements

This project was primarily funded through a U.S. Environmental Protection Agency (EPA) wetland protection grant to the Montana Department of Environmental Quality (DEQ), as authorized by Section 104(b)(3) of the Clean Water Act. Lynda Saul at DEQ and Steve Potts at EPA have been instrumental in administering this program, and they deserve special thanks for their continued support of the Montana Natural Heritage Program's wetland inventory initiative.

Many individuals provided information about high quality wetlands in the study area. Special thanks to Burt Williams (The Nature Conservancy), Chris Phelps (Montana Land Reliance), Jay Parks (Bureau of Land Management), Jim Hanson (Montana Department of Fish, Wildlife & Parks), Rich Johnson (U.S. Fish and Wildlife Service), and Jim Sparks, Kim Reid, and Pat Pierson (U.S.

Forest Service). Thanks also to the private landowners who allowed us to inventory wetlands on their property.

Heritage Program staff were essential to the completion of this project. Thanks to Martin Miller and Terrie Kenney for data processing, Cedron Jones and Whitney Weber for maps and GIS manipulations, Paul Hendricks for zoological surveys, John Carlson for rare animal information, Bonnie Heidel for botanical inventories and rare plant information, Steve Cooper and Jack Greenlee for field work, and Cathie Jean and Joy Lewis for editing. Thanks to Katrina Dixon (Scheuerman) at NRIS for help with final report formatting and production. Thanks also to Joe Elliott for identifying moss specimens and Peter Lesica for help with field surveys.

Table of Contents

INTRODUCTION	1
STUDY AREA	2
Physical Setting	2
Vegetation.....	3
Cultural Characteristics and Land Use	5
METHODS	6
Site Identification and Selection	6
Data Collection.....	7
Data Management	7
Site and Community Ranking	8
Community Rarity Ranks (State and Global Ranks)	8
Community Viability Ranks	8
Site Ranks	8
Plant Community Classification	9
Nomenclature	10
RESULTS AND DISCUSSION	10
Wetland Status	10
Plant Communities.....	12
Forested Vegetation.....	12
Scrub-Shrub Vegetation	16
Emergent Vegetation	16
Aquatic Bed Vegetation.....	17
Plant Species of Special Concern	17
Animal Species of Special Concern	17
Wetland Conservation	19
A-ranked Wetlands	21
B-ranked Wetlands	21
C-ranked Wetlands	31
Wetlands Not Inventoried	32
How This Information Can Be Used	33
Future Needs.....	33
How To Request Additional Information	33
LITERATURE CITED	35

APPENDIX A – Global and State Rank Guidelines

APPENDIX B – Site Rank Criteria for Wetlands and Riparian Areas

**APPENDIX C – Common Wetland Communities Documented in the upper Yellowstone River
Watershed**

APPENDIX D – Site Descriptions

APPENDIX E – Watersheds with High Biological Diversity and Conservation Value